



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ERRATUM

To accompany the paper by R. A. Daly on "The Classification of Igneous Intrusive Bodies," *Journal of Geology*, Vol. XIII, p. 485.

NOTE.—Through oversight the "explanations" of the figures in the text were omitted.

EXPLANATIONS OF FIGURES

FIG. 1.—Diagrammatic map of a composite dike in Arran.

(After Judd, *Quarterly Journal of the Geological Society*, London, Vol. XLIX (1893), p. 545.)

- | | |
|----------------------------|------------------------|
| 1. Country rock (granite). | 2. Augite andesite. |
| 3. Quartz-felsite. | 4. Pitchstone-porphry. |

The andesite was intruded along a fissure in the granite; then the felsite and porphyry were in succession intruded along the middle plane of the andesite dike.

FIG. 2.—Section of the Moyie Sill, British Columbia, at the crossing of the Moyie River and International Boundary.

Illustrating a body which has the form and relations of a true sill, although the thickness is more than 2,500 feet.

FIG. 3.—Section of a composite sill in the island of Skye.

(After Harker, *Tertiary Igneous Rocks of Skye*, 1904, p. 204.) The stratified Lias was cut by the sill of basalt; a later sill, of granophyre, was intruded along the middle plane of the basic sill. The latter itself may have been double.

FIG. 4.—Section of a composite laccolith.

(After Harker, *op. cit.*, p. 209.) The black is basalt, the white, granophyre. The laccolith cuts heavily bedded lava flows. The maximum thickness of the laccolith is 150 feet.

FIG. 5.—Section of an interformational laccolith.

(After Weed and Pirsson, *Journal of Geology*, Vol. IV (1896), p. 412.) The floor of the porphyry laccolith (in black) is composed of Pre-Cambrian crystalline schists; the cover, of Palæozoic sediments. The length of the section represented is about ten miles.

FIG. 6.—Section of a typical volcanic neck.

(After Geikie, *Ancient Volcanoes of Great Britain*, Vol. II. p. 273.)

FIG. 7.—Diagrammatic map of a Tertiary "chonolith" of rhombenporphyry (crosses).

Cutting intensely folded Palæozoic sediments (broken lines) and Tertiary sandstones and conglomerates (white), occurring on the Kettle River, British Columbia. At its northern end the porphyry disappears under a late Tertiary lava-cap (stippled). The Tertiary sediments are tilted and faulted. The faulting was accompanied, or immediately followed, by the intrusion of the porphyry which field evidence shows to have been injected as a body with a highly irregular form.

FIG. 8.—Diagrammatic map and section of Ascutney Mountain, Vermont.

Illustrating a composite stock, composed of successive intrusions, in stock or boss form, of diorite, syenite, and granite. A small boss of syenite (shown in black) cuts the diorite. These bodies cut crystalline schists, the attitude of which is indicated by the conventional symbol for strike and dip.

FIG. 9.—Map of part of the West Kootenay District, British Columbia.

Illustrating a small batholith of syenite (vertical lining) and the southern edge of the great composite batholith of the district. The latter is made up of the "Nelson granite" (thin, widely spaced, horizontal lines), cut by the younger "Rossland Alkali granite" (more closely set horizontal lines) and by the "Valhalla granite" (heavy horizontal lines). The maximum east-west width of the smaller batholith is fifteen miles.